


REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204 Arlington, VA 22202-4302 and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 07 December 1994		3. REPORT TYPE AND DATES COVERED Technical
4. TITLE AND SUBTITLE ON THE SYNTHESIS OF MIXED-METAL AND MIXED-PNICOGEN 13-15 RING SYSTEMS			5. FUNDING NUMBERS N00014-89-J-1545 R&T Project 4135008 Dr. Harold E. Guard	
6. AUTHOR(S)  R. L. Wells*, A. T. McPhail, P. S. White, M. S. Lube, and L J. Jones III				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Department of Chemistry Duke University Durham, NC 27708-0346			8. PERFORMING ORGANIZATION REPORT NUMBER  Technical Report No. DU/DC/TR-45	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)  Office of Naval Research 300 North Quincy Street Arlington, VA 22217-5000			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES  Accepted for Publication in <i>Phosphorus, Sulfur, and Silicon, and the Related Elements</i>				
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for Public Release Distribution Unlimited				
12b. DISTRIBUTION CODE				
13. ABSTRACT (Maximum 200 words)  Several pathways toward the synthesis of mixed-metal and mixed-pnicogen 13-15 ring systems have been explored. Equilibration reactions have proven successful toward the preparation of the latter compounds while most attempts to prepare mixed-metal compounds have resulted in unexpected and novel 13-15 species. These reactions and their products are discussed.				
19941213 039				
DTIC QUALITY INSPECTED 1				
14. SUBJECT TERMS 13-15 compounds, Mixed-Metal, Mixed-Pnicoge Ring Systems, Synthesis, Crystal Structure			15. NUMBER OF PAGES 4	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

OFFICE OF NAVAL RESEARCH

Grant N00014-89-J-1545

R&T Project 4135008

Dr. Harold E. Guard

Technical Report No. DU/DC/TR-45

**ON THE SYNTHESIS OF  
MIXED-METAL AND MIXED PNICOGEN 13-15 RING SYSTEMS**

by

R. L. Wells\*, A. T. McPhail, P. S. White, M. S. Lube, and L. J. Jones III

Accepted for Publication in *Phosphorus, Sulfur, and Silicon, and the Related Elements*

Duke University  
Department of Chemistry,  
P. M. Gross Chemical Laboratory  
Box 90346  
Durham, NC 27708-0346

07 December 1994

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
DTIC	Avail and/or Special
A-1	

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

## ON THE SYNTHESIS OF MIXED-METAL AND MIXED-PNICOGEN 13-15 RING SYSTEMS

RICHARD L. WELLS<sup>†</sup>, ANDREW T. McPHAIL<sup>†</sup>, PETER S. WHITE<sup>‡</sup>,  
MICHAEL S. LUBE<sup>†</sup>, and LEONIDAS J. JONES III<sup>†</sup>

<sup>†</sup>Department of Chemistry, Duke University, Durham, North Carolina 27708, U.S.A.

<sup>‡</sup>Department of Chemistry, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27514, U.S.A.

**Abstract** Several pathways toward the synthesis of mixed-metal and mixed-pnicogen 13-15 ring systems have been explored. Equilibration reactions have proven successful toward the preparation of the latter compounds while most attempts to prepare mixed-metal compounds have resulted in unexpected and novel 13-15 species. These reactions and their products are discussed.

### INTRODUCTION

Several compounds which have been structurally characterized by X-ray crystallography as dimeric in the solid-state have been shown to exhibit monomer-dimer equilibrium in solution.<sup>1,2</sup> Based on this phenomenon, reactions involving the equilibration of two dimers, each containing different group 15 elements, were attempted.

We have reported the facile conversion of dimeric compounds,  $[R_2ME(SiMe_3)_2]_2$ , to their respective mixed-bridge species,  $R_2\overline{ME(SiMe_3)_2M(R)_2}Cl$ , by reaction with two mole equivalents of  $R_2MCl$  ( $M = Ga, E = As^3$ ;  $M = In, E = P^4$  or  $As^5$ ) [eqn. 1]. In



addition, the conversion of a mixed-bridge species to its respective dimer can readily be accomplished via a salt elimination reaction using one mole equivalent of  $LiE(SiMe_3)_2$  ( $M = Ga, E = As^3$ ;  $M = In, E = P^6$ ;  $M = In, E = As^{7,8}$ ) [eqn. 2]. Each of these clean, high



yield reactions involved reagents which contained only the same group 13 and 15 elements and identical ligands. The success of these reactions prompted investigations into the synthesis of cyclic mixed-metal,  $\overline{M-E-M'-E}$ , and cyclic mixed-pnicogen,  $\overline{M-E-M-E'}$ , compounds.

EQUILIBRATION REACTIONSDimer Equilibrations

$[(\text{Me}_3\text{SiCH}_2)_2\text{InP}(\text{SiMe}_3)_2]_2$  (**1**) was found to exhibit monomer-dimer equilibrium at low concentrations as evidenced by the presence of both a doublet (monomer) and a triplet (dimer) in the  $^1\text{H}$  NMR spectrum arising from the virtual coupling of the trimethylsilyl protons with phosphorus.<sup>7</sup> Equimolar amounts of **1** and  $[(\text{Me}_3\text{SiCH}_2)_2\text{InAs}(\text{SiMe}_3)_2]_2$  (**2**) were dissolved in benzene and allowed to stir for 24 h at room temperature. Removing the volatiles and recrystallizing the solid residue from cold pentane yielded crystalline  $[(\text{Me}_3\text{SiCH}_2)_2\text{In}(\text{As/P})(\text{SiMe}_3)_2]_2$ . An X-ray crystallographic analysis of these crystals revealed a 50:50 occupancy for As and P at the pnictogen sites. The electron ionization (20 eV) mass spectrum showed a fragmentation pattern indicative of



$(\text{Me}_3\text{SiCH}_2)_2\overline{\text{InAs}(\text{SiMe}_3)_2\text{In}(\text{CH}_2\text{SiMe}_3)_2\text{P}(\text{SiMe}_3)_2}$  (**3**) and no peaks above  $m/z$  510 consistent with either **1** or **2**.<sup>9</sup> Clusters assigned to specific ions in the mass spectrum show appropriate isotopic patterns as calculated for the atoms present.

When  $[\text{I}_2\text{GaP}(\text{SiMe}_3)_2]_2$  (**4**) and  $[\text{I}_2\text{GaAs}(\text{SiMe}_3)_2]_2$  (**5**), both of which exhibit monomer-dimer equilibria in solution, were combined in a 1:1 mole ratio in toluene, the crystalline compound  $[\text{I}_2\text{Ga}(\text{As/P})(\text{SiMe}_3)_2]_2$  was isolated after 4 days. Like **3**, these crystals exhibited a 50:50 occupancy for As and P at the pnictogen sites. Electron ionization mass spectrometry unequivocally identified the compound as

$\text{I}_2\overline{\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{I})_2\text{P}(\text{SiMe}_3)_2}$  (**6**). Compounds **3** and **6**, both of which are extremely air sensitive, are the first examples of heavier group 13 mixed-pnictogen compounds.

Dimer and Diorganometal Halide Equilibrations

Attempts to prepare mixed-metal mixed-bridges by the equilibration of a group 13-15 dimer with a diorganometal halide [see eqn. 1] in which the halide and the dimer contained two different group 13 metals did not yield the desired compounds, but rather, unusual ligand redistribution products. The reaction of **2** with  $\text{Ph}_2\text{GaCl}$  [eqn. 3] and  $[\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2]_2$  (**7**)<sup>10</sup> with  $(\text{Me}_3\text{SiCH}_2)_2\text{InCl}$  [eqn. 4] gave the same mixed-ligand




## ON THE SYNTHESIS OF MIXED-METAL AND

indium-arsenic dimer  $[\text{Ph}(\text{Me}_3\text{SiCH}_2)\text{InAs}(\text{SiMe}_3)_2]_2$  (**8**).<sup>11</sup> Subsequently, compound **8** was isolated in a 49.0% yield by allowing **2** to react with 2 mole equivalents of  $\text{Ph}_3\text{Ga}$ , a yield which is much higher than those of the previous two reactions (8.7% and 7.8%, respectively). Compound **8** was found to reside in the trans configuration in the solid-state, however, the  $^1\text{H}$  NMR spectrum was consistent with the existence of equimolar amounts of the cis and trans isomers in solution at room temperature. Although it is not a mixed-metal compound, **8** was the first heavier group 13-15 compound to be fully characterized which contained two different organic substituents at the metal site. A reaction similar to that shown in eqn. 3, involving **1** and  $\text{Ph}_2\text{GaCl}$ , produced the halogen exchange product  $[\text{Cl}(\text{Me}_3\text{SiCH}_2)\text{InP}(\text{SiMe}_3)_2]_2$  rather than a phenyl exchange product.<sup>11</sup>

In an attempt to prevent the formation of a unique ligand exchange product, **2** was allowed to react with 2 mole equivalents of  $(\text{Me}_3\text{SiCH}_2)_2\text{GaCl}$  in toluene. The resulting viscous red oil was thermally unstable above  $-78^\circ\text{C}$ .  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra indicated that neither of the starting materials nor the dimer  $[(\text{Me}_3\text{SiCH}_2)_2\text{GaAs}(\text{SiMe}_3)_2]_2$  were present.

### SALT ELIMINATION REACTIONS

Reactions of gallium- and indium-arsenic mixed-bridging compounds with the lithium phosphides  $\text{LiP}(\text{SiMe}_3)_2$  and  $\text{LiPPh}_2$  were performed in an attempt to prepare compounds containing a four-membered ring incorporating two group 13 centers bridged by one arsenic and one phosphorus atom [eqn. 5]. When  $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Cl}$  (**9**)<sup>10</sup>



and  $\text{LiP}(\text{SiMe}_3)_2$  were combined in benzene at room temperature,  $\text{LiCl}$  and the unexpected ligand redistribution product  $\text{Ph}_3\text{Ga}\cdot\text{As}(\text{SiMe}_3)_3$  (**10**) (29.2% yield based on arsenic) were isolated.<sup>12</sup> Compound **10** was later prepared in a quantitative yield by the reaction of  $\text{Ph}_3\text{Ga}$  with  $\text{As}(\text{SiMe}_3)_3$ .

Upon the addition of a toluene solution of **9** to a 1 mole equivalent toluene suspension of  $\text{LiPPh}_2$ , the solid yellow phosphide appeared to react immediately. After stirring overnight at room temperature, a fine white precipitate was observed in the colorless solution. Filtering the solution separated  $\text{LiCl}$  from the liquor which, after removing the volatiles, yielded a toluene soluble white powder.  $^1\text{H}$  NMR and a partial elemental analysis were consistent with a compound of formula

$\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{PPh}_2$  [Anal. Calcd. (Found) for  $\text{C}_{42}\text{H}_{48}\text{AsGa}_2\text{PSi}_2$ : C, 59.05 (58.83); H, 5.66 (5.73)]. The  $^1\text{H}$  NMR spectrum consists of a singlet at  $\delta$  0.21

and a series of multiplets in the phenyl region, several of which appear to be further split due to virtual coupling to phosphorus. In a manner similar to that related above, the reaction of  $(\text{Me}_3\text{SiCH}_2)_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{CH}_2\text{SiMe}_3)_2\text{Cl}$  with  $\text{LiPPh}_2$  produced  $\text{LiCl}$  and a toluene soluble powder. Elemental analysis found an As:P ratio slightly greater than 1. Several small crystals were isolated and determined crystallographically to be  $[(\text{Me}_3\text{SiCH}_2)_2\text{GaAs}(\text{SiMe}_3)_2]_2$  (**11**).<sup>13</sup> The presence of this dimer would explain the deviation from unity of the As:P ratio, however, the sample consisted mainly of a powder rather than dimer **11** which readily forms large colorless crystals.

**Acknowledgement** Financial support for this work by the Office of Naval Research and the Lord Foundation of North Carolina is gratefully appreciated.

#### REFERENCES

1. R. L. Wells, M. F. Self, A. T. McPhail, S. R. Aubuchon, J. P. Jasinski, and R. C. Woudenberg, Organometallics, **12**, 2832 (1993).
2. O. T. Beachley, J. P. Kopasz, H. Zhang, W. E. Hunter, and J. L. Atwood, Organomet. Chem., **325**, 69 (1987).
3. R. L. Wells, A. T. McPhail, J. W. Pasterczyk, and A. Alvanipour, Organometallics, **11**, 226 (1992).
4. R. L. Wells, A. T. McPhail, and M. F. Self, Organometallics, **11**, 221 (1992).
5. R. L. Wells, L. J. Jones III, A. T. McPhail, and A. Alvanipour, Organometallics, **10**, 2345 (1991).
6. R. L. Wells, M. F. Self, unpublished results.
7. L. J. Jones III, Ph.D. Dissertation, Duke University (1993).
8. L. J. Jones III, A. T. McPhail, and R. L. Wells, Organometallics, in press.
9. L. J. Jones III, A. T. McPhail, and R. L. Wells, Organometallics, **13**, 2504 (1994).
10. W. K. Holley, R. L. Wells, S. Shafieezad, A. T. McPhail, and C. G. Pitt, J. Organometallic Chem., **381**, 15 (1990).
11. R. L. Wells, A. T. McPhail, L. J. Jones III, and M. F. Self, J. Organomet. Chem., **449**, 85 (1993).
12. R. L. Wells, A. T. McPhail, L. J. Jones III, and M. F. Self, Organometallics, **11**, 2694 (1992).
13. R. L. Wells, J. W. Pasterczyk, A. T. McPhail, J. D. Johansen and A. Alvanipour, J. Organomet. Chem., **407**, 17 (1991).

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (1)\*  
Chemistry Division, ONR 331  
800 North Quincy Street  
Arlington, Virginia 22217-5660

Dr. Richard W. Drisko (1)  
Naval Facilities & Engineering  
Service Center  
Code L52  
Port Hueneme, CA 93043

Defense Technical Information  
Center (2)  
Building 5, Cameron Station  
Alexandria, VA 22314

Dr. Eugene C. Fischer (1)  
Code 2840  
Naval Surface Warfare Center  
Carderock Division Detachment  
Annapolis, MD 21402-1198

Dr. James S. Murday (1)  
Chemistry Division, Code 6100  
Naval Research Laboratory  
Washington, D.C. 20375-5320

Dr. Bernard E. Douda (1)  
Crane Division  
Naval Surface Warfare Center  
Crane, Indiana 47522-5000

Dr. John Fischer, Director (1)  
Chemistry Division, C0235  
Naval Air Weapons Center  
Weapons Division  
China Lake, CA 93555-6001

Dr. Peter Seligman (1)  
Naval Command, Control and  
Ocean Surveillance Center  
RDT&E Division  
San Diego, CA 92152-5000

\* Number of copies to forward